


# YOUR DIABETIC GASTROPARESIS DOCTOR DISCUSSION GUIDE



## Making the Most of Your Doctor Visit

Use this discussion guide to help you speak with a doctor about your symptoms. This can help your doctor review your current treatment plan and determine the appropriate next steps for you.

**On a scale of 1 to 5, rate how well your current treatment for diabetic gastroparesis is working for you.**



**1**

Not working at all



**2**



**3**



**4**



**5**

My symptoms are greatly reduced



I am not currently taking any treatment

**Which diabetic gastroparesis symptoms are most common for you? Check all that apply.**



Nausea



Vomiting



Bloating



Stomach pain



Feeling full after eating only a small amount



**Other** (please specify) \_\_\_\_\_

**On a scale of 1 to 10, how much have your diabetic gastroparesis symptoms affected your daily activities?**



**1-3**

I live my life the way I want to on most days



**4-6**

My symptoms flare up weekly



**7-9**

My symptoms dictate my life most days



**10**

I am completely at the mercy of my symptoms

Please see complete Important Safety Information on pages 4-5, including Boxed Warning. Also see complete [Prescribing Information](#), [Medication Guide](#), and [Instructions for Use](#).

 **Gimoti**<sup>®</sup>  
(metoclopramide)  
nasal spray

## What aspects of your life have changed the most because of your symptoms? Check all that apply.

### Relationships

I can't spend as much time with my family and friends as I want to

### Work and career

I have to reduce my work hours or quit working

### Physical health

I can't be as active as I want to be

### Mental health

I often feel hopeless because of my symptoms

Other (please specify) \_\_\_\_\_

## Do you have difficulty taking or keeping down pills because of your diabetic gastroparesis symptoms?

Yes

No

### What is the most important information I should know about GIMOTI?

GIMOTI can cause serious side effects, including: **TARDIVE DYSKINESIA: Abnormal muscle movements, mostly of the face or tongue muscles. You cannot control these movements, and they may not go away even after stopping GIMOTI. Your chances of getting tardive dyskinesia increase**

- **The longer you take metoclopramide and the more metoclopramide you take. You should not take GIMOTI for more than 8 weeks at a time, and you should not take products containing metoclopramide (including GIMOTI) for more than 12 weeks at a time.**
- **If you are older, especially if you are an older woman (e.g., age 65 years and older)**
- **If you have diabetes**

---

**Call your healthcare professional right away if you get movements you cannot stop or control, such as lip smacking, chewing, or puckering up your mouth; frowning or scowling; sticking out your tongue; blinking and moving your eyes; shaking of your arms and legs.**

---

Please see complete Important Safety Information on pages 4-5, including Boxed Warning. Also see complete [Prescribing Information](#), [Medication Guide](#), and [Instructions for Use](#).

 **Gimoti**<sup>®</sup>  
(metoclopramide)  
nasal spray

# ADVOCATE FOR YOUR HEALTH



Many people with diabetic gastroparesis think they have to settle for living with symptoms like nausea, vomiting, and stomach pain. **If diet and lifestyle changes or your current medications aren't working for you, there are other options.**

## Tips for talking to your doctor:



**Be prepared.** Go in ready to talk about which symptoms are the most bothersome, what in your treatment plan is not working, and your goals for symptom management.



**Be open.** The more you tell your doctor about what's going on, the more he or she will be able to help.



**Be clear and specific.** For example, instead of saying, "My stomach hurts when I eat," talk about how severe your symptoms are and how they affect you. You could say, "My stomach hurts so badly after a small meal that I sometimes won't eat to avoid feeling that way."



**Be your own best advocate.** Don't be afraid to share that you want to make a change or to ask for what you need. You may also consider bringing a family member or friend for additional support. You deserve to get your symptoms under control!

## Questions to ask your doctor when considering Gimoti® (metoclopramide) nasal spray:

- Since my current treatment plan isn't working, could it help to make a change?
- Could GIMOTI be the right choice for me?
- How is GIMOTI different from other treatments?
- How do I take GIMOTI?
- What are the possible side effects of GIMOTI?
- How will I know if GIMOTI is working? What benefits can I expect?

Please see complete Important Safety Information on pages 4-5, including Boxed Warning. Also see complete [Prescribing Information](#), [Medication Guide](#), and [Instructions for Use](#).

 **Gimoti**<sup>®</sup>  
(metoclopramide)  
nasal spray

## INDICATION AND IMPORTANT SAFETY INFORMATION

### What is Gimoti® (metoclopramide) nasal spray?

GIMOTI is a prescription medicine used 30 minutes before meals and at bedtime for 2 to 8 weeks to relieve symptoms of slow stomach emptying in adults with diabetes. Avoid treatment with metoclopramide (all dosage forms and routes of administration) for longer than 12 weeks. GIMOTI is not recommended for use in children under age 18.

### Important facts about GIMOTI

This is a summary of important information you need to know about GIMOTI. This information does not take the place of talking with your healthcare professional about your medical condition or your treatment.

### What is the most important information I should know about GIMOTI?

**GIMOTI can cause serious side effects, including: TARDIVE DYSKINESIA: Abnormal muscle movements, mostly of the face or tongue muscles. You cannot control these movements, and they may not go away even after stopping GIMOTI. Your chances of getting tardive dyskinesia increase**

- **The longer you take metoclopramide and the more metoclopramide you take. You should not take GIMOTI for more than 8 weeks at a time, and you should not take products containing metoclopramide (including GIMOTI) for more than 12 weeks at a time.**
- **If you are older, especially if you are an older woman (e.g., age 65 years and older)**
- **If you have diabetes**

Call your healthcare professional right away if you get movements you cannot stop or control, such as lip smacking, chewing, or puckering up your mouth; frowning or scowling; sticking out your tongue; blinking and moving your eyes; shaking of your arms and legs.

### Do not use GIMOTI if you

- Have a history of tardive dyskinesia or have a problem controlling your muscles and movements after taking GIMOTI or a medicine that works like GIMOTI
- Have stomach or intestinal problems that could get worse with GIMOTI, such as bleeding, blockage, or a tear in the stomach or bowel wall

- Have a type of tumor that can cause high blood pressure, such as pheochromocytoma
- Have epilepsy (seizures)
- Are allergic to metoclopramide. Stop taking GIMOTI right away and get emergency help if you have any of these symptoms:
  - swelling of your tongue, throat, lips, eyes, or face
  - trouble swallowing or breathing
  - skin rash, hives, sores in your mouth, or skin blisters

**Before starting GIMOTI**, tell your healthcare professional about all your medical conditions, especially if you have

- Problems controlling your muscle movements after taking any medicine
- Parkinson's disease
- Pheochromocytoma
- Kidney or liver disease
- Depression or mental illness
- High blood pressure

Also tell your healthcare professional if you are pregnant or plan to become pregnant, are breastfeeding or plan to breastfeed, or drink alcohol.

Tell your healthcare professional about all the medicines you take, including prescription drugs and over-the-counter medicines, vitamins, and herbal supplements.

Speak with your healthcare professional before you start or stop any other medicines.

### Especially tell your healthcare professional if you take

- Another medicine that contains metoclopramide, such as REGLAN® tablets
- Medicine for Parkinson's disease
- Blood pressure medicine
- Medicine for depression, especially a monoamine oxidase inhibitor (MAOI)
- Antipsychotic medicine used to treat mental illness, such as schizophrenia
- Insulin
- Medicines that can make you sleepy, such as anxiety medicines, sleep medicines, and narcotics



**What should I avoid while taking GIMOTI?**

- Do not drink alcohol while taking GIMOTI
- GIMOTI may cause sleepiness or dizziness. Do not drive, operate machinery, or do potentially dangerous activities until you know how GIMOTI affects you

**What are other possible side effects of GIMOTI?**

- Other changes in muscle control and movement, such as:
  - uncontrolled spasms of your face and neck muscles, or muscles of your body, arms, and legs (dystonia)
  - parkinsonism – slight shaking, body stiffness, and trouble moving or keeping your balance
  - being unable to sit still or feeling that you need to move your hands, feet, or body (akathisia)
- Neuroleptic Malignant Syndrome (NMS) – a very rare but very serious condition. NMS can lead to death and must be treated in a hospital
- Depression, thoughts about suicide, and suicide
- High blood pressure
- Too much body water
- Increased prolactin

**Call your healthcare professional and get medical help right away if you**

- Feel depressed or have thoughts about hurting or killing yourself
- Have high fever, stiff muscles, problems thinking, very fast or uneven heartbeat, and/or increased sweating
- Have muscle movements that you cannot stop or control
- Have muscle movements that are new or unusual

**The most common side effects of GIMOTI include**

- Unpleasant taste after dosing
- Headache
- Tiredness

These are not all the possible side effects of GIMOTI. Ask your healthcare professional for medical advice about side effects.

You may report side effects related to Evoke Pharma products by calling 1-833-4-GIMOTI (1-833-444-6684) or emailing [GIMOTImedinfo@evokepharma.com](mailto:GIMOTImedinfo@evokepharma.com). If you prefer to report these to the FDA, either visit [www.FDA.gov/medwatch](http://www.FDA.gov/medwatch) or call 1-800-FDA-1088.

This information should not take the place of you talking with your doctor or healthcare professional. If you have any questions about your condition, or if you would like more information about GIMOTI, talk to your doctor or pharmacist. Only you and your healthcare professional can decide if GIMOTI is right for you.

**Please see complete [Prescribing Information](#), including [Boxed Warning](#), [Medication Guide](#), and [Instructions for Use](#).**